

2021

## PHYSICS — HONOURS

## Third Paper

Full Marks : 100

*The figures in the margin indicate full marks.**Candidates are required to give their answers in their own words as far as practicable.***Question no. 1** is compulsory. Answer **any eight** questions taking **four** from each **unit**.1. Answer **any ten** questions :

2×10

- (a) What is a multiplexer? Draw its 4 to 1 circuit diagram.
- (b) The modulation index of an FM wave is 10 and the highest modulation frequency is 20 KHz. Find the minimum bandwidth required for the detection of the FM wave.
- (c) Build half adder with minimum number of NAND gates.
- (d) Write the integral form of Faraday's law and hence show that  $\nabla \times \vec{E} = -\frac{\partial \vec{B}}{\partial t}$ .
- (e) An AC circuit connected to a 220V, 50 Hz supply contains a 20H coil of resistance 100  $\Omega$ , connected in series with 1  $\mu$ F capacitor. Calculate the power factor of the circuit.
- (f) What is series resonant circuit and why is it called acceptor circuit?
- (g) The potential in a region is expressed as  $\phi = 2\beta(x^2 + y^3 + z)$ . Calculate the charge density in this region.
- (h) A pure dipole 'p' is placed at the origin, pointing in the z-direction. Find the force on a point charge q at (a, 0, 0).
- (i) Define dielectric susceptibility of a medium. What is non-linear dielectric?
- (j) A single slit, illuminated by a red light of  $6500 \text{ \AA}$ , gives first-order Fraunhofer diffraction minima that have angular distances of  $\theta = 5^\circ$  from the optic axis. How wide is the slit?
- (k) What is meant by 'missing order' in a double slit diffraction pattern?
- (l) State Brewster's law.

## Unit - 5

2. (a) Define the quantity *Common Mode Rejection Ratio* ( $\rho$ ) for a Differential Amplifier.
- (b) For a Differential Amplifier if  $v_1$  and  $v_2$  are the input signals, then show that the output signal  $v_o$  is given by :

$$v_o = A_d v_d (1 + v_c / \rho v_d),$$

where  $v_d$  is the difference voltage and  $v_c$  is the common mode signal and  $A_d$  is the gain of the differential Amplifier.

Please Turn Over

(c) Draw an analog circuit using OPAMP to solve the following simultaneous equations :

(i)  $5x + 2y = 12$  (ii)  $2x + 3y = 6$

(d) Show that the negative feedback improves the stability of an amplifier.

1+3+3+3

3. (a) Why negative resistance is provided in an oscillator?

(b) Draw the circuit diagram of a tuned-collector oscillator and write the expression for the frequency of oscillation.

(c) State why *crystal oscillators* are superior for high frequency stability over *LC oscillators*.

(d) Draw the circuit of an astable multivibrator using transistors. Draw the waveforms at the collector of any one transistor.

2+(2+1)+2+(2+1)


4. (a) Draw a clocked S-R flip-flop circuit using two-input NAND gates and explain its operation with proper truth table.

(b) Distinguish between combinational and sequential logic circuits.

(c) Draw the circuit diagram of a 4-bit ripple counter using 4JK flip-flops (FF). Explain its working.

(2+3)+2+3

5. (a) Find the force on a square loop placed as shown in the following figure near an infinite straight wire. Both the loop and the wire carry steady current  $I$ .



(b) Show that the magnetic field at a large distance ( $\vec{r}$ ) due to a small current loop having magnetic moment  $\vec{m}$  is given by

$$\vec{B}(\vec{r}) = \frac{\mu_0}{4\pi} \left[ -\frac{\vec{m}}{r^3} + \frac{3(\vec{m} \cdot \vec{r})\vec{r}}{r^5} \right]$$

(c) Magnetic vector potential is given by  $\vec{A} = e^{-x} \sin y \hat{i} + (1 + \cos y) \hat{j}$ . Calculate the magnetic induction.

3+4+3

6. (a) Show that the energy of a magnetic dipole in a magnetic field  $\vec{B}$  is given by  $U = -\vec{m} \cdot \vec{B}$ .

(b) Plot the variation of relative permeability of a magnetic material with magnetic field intensity during magnetization. Explain the nature of plot. How can a material be demagnetized from its initial state of magnetization? Explain with the help of hysteresis loop.

(c) Two magnetic media are separated by a plane interface. Show that the angles between the normal to the boundary and the  $\vec{B}$  fields on either side satisfy  $\mu_2 \tan \theta_1 = \mu_1 \tan \theta_2$  [symbols having usual meaning].

3+4+3


7. (a) A circular loop of wire, with radius  $R$ , lies in  $xy$ -plane, centered at the origin and carries a current  $I$  running counterclockwise as viewed from the positive  $z$  axis.
- What is the magnetic dipole moment?
  - What is its (approximate) magnetic field at points far from the origin?
- (b) Determine the self inductance of a co-axial cable.
- (c) Show the equivalent inductance of two inductances joined in parallel is given by

$$L_{eqv} = \frac{L_1 L_2 - M^2}{L_1 + L_2 \mp 2M} \quad (1+2)+3+4$$

### Unit - 6

8. (a) Two charges of charge  $+q$  and  $-q$  are situated at the location  $(a, 0, 0)$  and  $(-a, 0, 0)$ . If a large conducting plane is placed above at  $z = a$ , then calculate the net force on  $+q$  charge.
- (b) Show that the dipole moment of a charge distribution is independent of the choice of the origin if the total charge vanishes.
- (c) Three charges are situated at the corners of a square (side  $a$ ) as shown in the figure below. How much work does it take to bring another charge,  $+q$ , from far away and place it in the fourth corner?

4+3+3



9. (a) Find the electric field at a distance  $z$  above the midpoint of a straight line segment of length  $2L$ , which carries a uniform line charge  $\lambda$ . Show that, for points far from the line, the line 'looks' like a point charge  $q = 2\lambda L$ .
- (b) Which one of the following is an impossible electrostatic field?
- $\vec{E} = k[xy\hat{x} + 2yz\hat{y} + 3xz\hat{z}]$
  - $\vec{E} = k[y^2\hat{x} + (2xy + z^2)\hat{y} + 2yz\hat{z}]$
- (c) Spherical charge distribution has been expressed as :

$$\begin{aligned} \rho &= \rho_0(1 - r^2/a^2) & \text{for } r \leq a \\ &= 0 & \text{for } r > 0 \end{aligned}$$

Find the electric field intensity and potential inside ( $r < a$ ) the charge distribution. (3+1)+2+4

**Please Turn Over**

10. (a) Find the general solution to Laplace's equation in spherical coordinates, for the case where the potential,  $V$  depends on  $r$  only.
- (b) Prove, by the method of boundary condition, that the field inside a spherical cavity in an isotropic dielectric is given by :  $E_i = E + (P / 3\epsilon_0)$ , where the symbols have their usual meaning.
- (c) Show that the electric field produced by a polarized dielectric can be given by the contributions from a bound charge density  $\sigma = \vec{P} \cdot \hat{n}$  and a volume charge density  $\rho = -\vec{\nabla} \cdot \vec{P}$ . 2+4+4
11. (a) Show that, in two dimensions the shape of the fringes produced in Young's experiment is hyperbolic.
- (b) Cite one example / experiment for the each following type of interferometers :
- (i) Wavefront-splitting interferometer (ii) Amplitude-splitting interferometer.
- (c) A soap film of refractive index 1.33 and of thickness  $1.4 \mu\text{m}$  is illuminated by white light falling at an angle of  $60^\circ$ . The light reflected by it contains a dark band corresponding to wavelength of  $5000\text{\AA}$ . Determine the order of interference dark band.
- (d) Why a broad source of light is necessary for observing colours in thin film? 3+2+3+2
12. (a) Derive an expression for the intensity of Fraunhofer diffraction pattern for a double slit.
- (b) Starting from the grating equation, find an expression describing the angular spread for a small range of wavelengths,  $\Delta\theta / \Delta\lambda$ , i.e., the dispersive power  $D$ . Then compute the dispersive power in the first and second orders for a grating with 1500 grooves per inch operating in the visible region.
- (c) Consider a grating with slit of width  $a = 0.001 \text{ mm}$ , separated by a distance of  $0.002 \text{ mm}$ . How many orders would be visible at  $\lambda = 500 \text{ nm}$ ? 4+(1+2)+3
13. (a) What is optical activity? Give Fresnel's explanation of rotation of plane of polarisation by an optically active substance.
- (b) An unpolarized beam strikes the surface of a pond. Find out the angle of incidence, so that the reflected beam will be completely polarized with its E-field perpendicular to the plane of incidence. Given : Refractive index of the pond water : 1.33. Derive the formula you use.
- (c) Explain the phenomenon of double refraction in a uniaxial crystal by applying Huygen's theory. 4+(2+2)+2
-