

2021

## PHYSICS—HONOURS

Paper : CC-5

Full Marks : 50

*The figures in the margin indicate full marks.  
Candidates are required to give their answers in their own words  
as far as practicable.*

Answer **question no. 1** and **any four** from the rest.

1. Answer **any five** questions:

2×5

- Find the average value of  $\sin x + \sin^2 x$  for  $0 \leq x \leq 2\pi$ .
- Using generating function, show that  $p_l(-x) = (-1)^l p_l(x)$ , where  $P_l(x)$  is Legendre polynomial of order  $l$ .
- Find the solution of the equation  $x^2 \frac{d^2 y}{dx^2} + px \frac{dy}{dx} + qy = 0$ , where  $p$  and  $q$  are constants.
- For the Poisson's distribution  $P_n = \frac{\mu^n}{n!} e^{-\mu}$ . Find the standard deviation for the distribution.  $\mu$  is constant.

**Or**, [syllabus 2018-2019]

If the Lagrangian is invariant under a rigid translation, the momentum of the system is conserved.


- If  $f(x) \rightarrow 0$  for  $x \rightarrow \pm\infty$ , find the Fourier transform of  $\frac{df}{dx}$ .

**Or**, [syllabus 2018-2019]

Show that if the Hamiltonian is not explicitly time dependent then energy is conserved (assuming Hamiltonian equals energy).


- Show that  $x = 1$  is a regular singular point of the Legendre differential equation.
- Show that, for any  $p > 0$ ,  $\Gamma(p + 1) = p\Gamma(p)$ .

2. Consider the function shown in figure



- Find its Fourier transform  $g(\omega)$ . Sketch  $g(\omega)$  vs  $\omega$ .
- Show, using Parseval's identity  $\int_0^\infty \frac{\sin^2 \omega}{\omega^2} d\omega = \frac{\pi}{2}$ .
- Show that Fourier transform of  $f(t - t_0) = e^{-i\omega t_0} g(\omega)$ .

**Please Turn Over**



Using (a) and (c), find Fourier transform of  $h(t)$ .

(2+1)+3+2+2

**Or**, [Syllabus 2018-2019]

- (a) Find the path followed by a light ray if the index of refraction in polar coordinate is proportional to  $r^{-2}$ .  
 (b) Find the equation of motion of a particle moving along  $x$ -axis under a potential energy  $V = \frac{1}{2}kx^2$ , by constructing the Lagrangian. Construct the Hamiltonian for the system and argue that it is a conservative system.

5+5

3. (a) Calculate, using gamma function  $\int_1^\infty \frac{(\ln x)^3}{x^2(x-1)} dx$  (you can assume  $\sum_{r=1}^\infty \frac{1}{r^4} = \frac{\pi^4}{90}$ ).

- (b) Prove that for positive integers  $m$  and  $n$ ,  $\beta(m, n) = \frac{(n-1)!}{m.(m+1) \dots (m+n-1)}$ .

Hence show that  $1.3.5 \dots (2n-1) = \frac{2^n \Gamma(n+\frac{1}{2})}{\sqrt{\pi}}$ .

5+(3+2)

4. (a) Given  $f(x) = x$  for  $0 < x < 1$ , sketch even function corresponding to this function with period 2. Find the Fourier series for this even function.

- (b) Using above expansion, find the value of  $\sum_{n=1}^\infty \frac{1}{(2n-1)^2}$ .

- (c) Sketch the odd function of period 2 corresponding to the above given function.

5+3+2

5. (a) From the generating function of Hermite polynomial  $H_n(x)$ ,  $e^{2xt-t^2} = \sum_{n=0}^\infty \frac{1}{n!} H_n(x)$

Show that  $H'_n(x) = 2n H_{n-1}(x)$ .

- (b)  $H_n(x)$  are orthogonal polynomials in the domain  $-\infty < x < \infty$ . Suppose  $H_2(x) = a + bx^2$ . Find  $a$  and  $b$  using orthogonality property of  $H_n(x)$ ; given  $H_0(x) = 1$ ,  $H_1(x) = 2x$  and coefficient of highest power of  $x$  in  $H_n(x)$  is  $2^n$ .

- (c) Solve  $x^2 y'' - 6y = 0$  using Frobenius method around  $x = 0$ .

2+4+4

6. (a) For a binomial distribution with  $n$  trials, if  $p$  is the probability of success and  $q$  is that of failure, then show that the mean and variance of the distribution are respectively  $np$  and  $npq$ .

- (b) Solve  $\frac{\partial^2 y}{\partial x^2} = c \frac{\partial y}{\partial t}$  using Fourier transform.

- (c) A random variable  $x$  has the density function  $f(x) = \begin{cases} cx & 0 \leq x \leq 2 \\ 0 & \text{otherwise} \end{cases}$

find (i) the constant  $c$ .

(ii) the probability that  $x > 1$ .

(2+2)+3+(1+2)

**Or,** [Syllabus 2018-2019]

- (a) Derive the Euler-Lagrange equation from the Principle of Least Action. Show clearly how the variation of paths is implemented in your derivation in terms of ordinary partial derivative with respect to a parameter.

- (b) A bead slides on a frictionless wire in the shape of a cycloid described by the equations

$$x = a(\theta - \sin\theta) \quad y = a(1 + \cos\theta)$$

where  $(x, y)$  are cartesian coordinates,  $a$  is a constant and  $0 \leq \theta \leq 2\pi$ . Write down (i) the Lagrangian and (ii) the equation of motion in terms of  $\theta$ . Indicate the generalized coordinate  $\theta$  by drawing a figure.

5+(2+2+1)

7. (a) The heat equation in 2 dimensional cartesian coordinates is given by  $\frac{\partial^2 T}{\partial x^2} + \frac{\partial^2 T}{\partial y^2} = \frac{1}{K} \frac{\partial T}{\partial t}$

where  $T$  is the temperature function and  $K$  is a constant. Solve this equation for steady state using the method of separation of variables.

- (b) Find a solution  $U(x, t)$  of the boundary-value problem  $\frac{\partial U}{\partial t} = 3 \frac{\partial^2 U}{\partial x^2} \quad t > 0, \quad 0 < x < 2$

The boundary conditions are

$$U(0, t) = 0, \quad U(2, t) = 0$$

$$U(x, 0) = x \quad 0 < x < 2$$

$$\left[ \text{Given } x = \sum_{n=1}^{\infty} -\frac{4}{n\pi} (1)^n \sin \frac{n\pi x}{2}; 0 < x < 2 \right]$$

3+7